

SALBIN INHALER 100 micrograms

Summary of Product Characteristics | AMB HK ENTERPRISES (PVT) LTD

1. Name of the medicinal product

SALBIN INHALER 100 micrograms

2. Qualitative and quantitative composition

SALBIN INHALER is a pressurized metered-dose inhaler delivering 100 micrograms of salbutamol (as Salbutamol Sulfate BP) per actuation. SALBIN INHALER contains a new propellant (HFA 134a) and does not contain any chlorofluorocarbons.

3. Pharmaceutical form

Aerosol

4. Clinical particulars

4.1 Therapeutic indications

SALBIN INHALER is indicated in adults, adolescents and children aged 4 to 11 years.

SALBIN INHALER provides short-acting (4 to 6 hour) bronchodilation with fast onset (within 5 minutes) in reversible airways obstruction.

It is particularly suitable for the relief and prevention of asthma symptoms. It should be used to relieve symptoms when they occur, and to prevent them in those circumstances recognized by the patient to precipitate an asthma attack (e.g. before exercise or unavoidable allergen exposure).

SALBIN INHALER is particularly valuable as relief medication in mild, moderate or severe asthma, provided that reliance on it does not delay the introduction and use of regular inhaled corticosteroid therapy.

4.2 Posology and method of administration

SALBIN INHALER is for oral inhalation use only. SALBIN INHALER may be used with a Volumetric spacer device by patients who find it difficult to synchronize aerosol actuation with inspiration of breath.

Adults (including the elderly):

For the relief of acute asthma symptoms including bronchospasm, one inhalation (100 micrograms) may be administered as a single minimum starting dose. This may be increased to two inhalations if necessary. To prevent allergen- or exercise-induced symptoms, two inhalations should be taken 10-15 minutes before challenge.

For chronic therapy, two inhalations up to four times a day.

Paediatric Population

Relief of acute bronchospasm

The usual dosage for children under the age of 12 years: one inhalation (100 micrograms). The dose may be increased to two inhalations if required.

Children aged 12 years and over: dose as per adult population.

Prevention of allergen or exercise-induced bronchospasm

The usual dosage for children under the age of 12 years: one inhalation (100 micrograms) before challenge or exertion.

The dose may be increased to two inhalations if required.

Children aged 12 years and over: dose as per adult population.

Chronic therapy

The usual dosage for children under the age of 12 years: up to two inhalations 4 times daily.

Children aged 12 years and over: dose as per adult population.

The Babyhaler spacer device may be used to facilitate administration to children under 5 years of age.

On-demand use of SALBIN INHALER should not exceed 8 inhalations in any 24 hours. Reliance on such frequent supplementary use, or a sudden increase in dose, indicates poorly controlled or deteriorating asthma

4.3 Contraindications

Hypersensitivity to the active substance or any of the excipients.

Non-i.v. formulations of salbutamol must not be used to arrest uncomplicated premature labour or threatened abortion.

4.4 Special warnings and precautions for use

Patients inhaler technique should be checked to make sure that aerosol actuation is synchronized with inspiration of breath for optimum delivery of drug to the lungs. Patients should be warned that they may experience a different taste upon inhalation compared to their previous inhaler.

Bronchodilators should not be the only or main treatment in patients with severe or unstable asthma. Severe asthma requires regular medical assessment, including lung-function testing, as patients are at risk of severe attacks and even

death. Physicians should consider using the maximum recommended dose of inhaled corticosteroid and/or oral corticosteroid therapy in these patients.

The dosage or frequency of administration should only be increased on medical advice. If a previously effective dose of inhaled salbutamol fails to give relief lasting at least three hours, the patient should be advised to seek medical advice. Patients who are prescribed regular anti-inflammatory therapy (e.g., inhaled corticosteroids) should be advised to continue taking their anti-inflammatory medication even when symptoms decrease, and they do not require SALBIN INHALER. Increasing use of bronchodilators, in particular short-acting inhaled beta₂-agonists, to relieve symptoms, indicates deterioration of asthma control, and patients should be warned to seek medical advice as soon as possible. The patient should be instructed to seek medical advice if short-acting relief bronchodilator treatment becomes less effective, or more inhalations than usual are required. In this situation the patient should be assessed and consideration given to the need for increased anti-inflammatory therapy (e.g. higher doses of inhaled corticosteroid or a course of oral corticosteroid). Overuse of short-acting beta-agonists may mask the progression of the underlying disease and contribute to deteriorating asthma control, leading to an increased risk of severe asthma exacerbations and mortality.

Patients who take more than twice a week “as needed” salbutamol, not counting prophylactic use prior to exercise, should be re-evaluated (i.e., daytime symptoms, night-time awakening, and activity limitation due to asthma) for proper treatment adjustment as these patients are at risk for overuse of salbutamol.

Severe exacerbations of asthma must be treated in the normal way.

Cardiovascular effects may be seen with sympathomimetic drugs, including salbutamol. There is some evidence from post-marketing data and published literature of rare occurrences of myocardial ischemia associated with salbutamol. Patients with underlying severe heart disease (e.g. ischemic heart disease, arrhythmia or severe heart failure) who are receiving salbutamol should be warned to seek medical advice if they experience chest pain or other symptoms of worsening heart disease. Attention should be paid to assessment of symptoms such as dyspnea and chest pain, as they may be of either respiratory or cardiac origin.

Salbutamol should be administered cautiously to patients with thyrotoxicosis.

Potentially serious hypokalemia may result from beta₂-agonist therapy, mainly from parenteral and nebulized administration. Particular caution is advised in acute severe asthma as this effect may be potentiated by hypoxia and by concomitant treatment with xanthine derivatives, steroids and diuretics. Serum potassium levels should be monitored in such situations.

As with other inhalation therapy, paradoxical bronchospasm may occur with an immediate increase in wheezing after dosing. This should be treated immediately with an alternative presentation or a different fast-acting inhaled bronchodilator. INHALER should be discontinued immediately, the patient assessed, and if necessary, a different fast-acting bronchodilator instituted for on-going use.

4.5 Interaction with other medicinal products and other forms of interaction

Salbutamol and non-selective β₁-blocking drugs such as propranolol, should not usually be prescribed together.

4.6 Fertility, pregnancy and lactation

Pregnancy

Studies in animals have shown reproductive toxicity. Safety in pregnant women has not been established. No controlled clinical trials with salbutamol have been conducted in pregnant women. Rare reports of various congenital anomalies following intrauterine exposure to salbutamol (including cleft palate, limb defects and cardiac disorders) have been received. Some of the mothers were taking multiple medications during their pregnancies. SALBIN INHALER should not be used during pregnancy unless clearly necessary.

Breast-feeding

As salbutamol is probably secreted in breast milk, its use in nursing mothers requires careful consideration. It is not known whether salbutamol has a harmful effect on the neonate, and so its use should be restricted to situations where it is felt that the expected benefit to the mother is likely to outweigh any potential risk to the neonate.

Fertility

There is no information on the effects of salbutamol on human fertility. There were no adverse effects on fertility in animals.

4.7 Effects on ability to drive and use machines

None reported.

4.8 Undesirable effects

Adverse events are listed below by system organ class and frequency. Frequencies are defined as: very common (≥ 1/10), common (≥ 1/100 to <1/10), uncommon (≥ 1/1000 to <1/100), rare (≥ 1/10,000 to <1/1000) and very rare (<1/10,000)

including isolated reports. Very common and common events were generally determined from clinical trial data. Rare, very rare and unknown events were generally determined from spontaneous data.

Immune system disorders

Very rare: Hypersensitivity reactions including angioedema, urticaria, bronchospasm, hypotension and collapse.

Metabolism and nutrition disorders

Rare: Hypokalemia.
Potentially serious hypokalemia may result from beta-2 agonist therapy.

Nervous system disorders

Common: Tremor, headache.

Very rare: Hyperactivity.

Cardiac disorders

Common: Tachycardia.

Uncommon: Palpitations.

Very rare: Cardiac arrhythmias (including atrial fibrillation, supraventricular tachycardia and extrasystoles).

Unknown: Myocardial ischemia* (see section 4.4)

Vascular disorders

Rare: Peripheral vasodilatation.

Respiratory, thoracic and mediastinal disorders

Very rare: Paradoxical bronchospasm.

Gastrointestinal disorders

Uncommon: Mouth and throat irritation.

Musculoskeletal and connective tissue disorders

Uncommon: Muscle cramps.

* reported spontaneously in post-marketing data therefore frequency regarded as unknown

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

4.9 Overdose

The most common signs and symptoms of overdose with salbutamol are transient beta agonist pharmacologically mediated events, **including tachycardia, tremor, hyperactivity and metabolic effects including hypokalemia.** Hypokalemia may occur following overdose with salbutamol. Serum potassium levels should be monitored. Lactic acidosis has been reported in association with high therapeutic doses as well as overdoses of short-acting beta-agonist therapy, therefore monitoring for elevated serum lactate and consequent metabolic acidosis (particularly if there is persistence or worsening of tachypnea despite resolution of other signs of bronchospasm such as wheezing) may be indicated in the setting of overdose.

5. Pharmacological properties

5.1 Pharmacodynamics properties

Pharmacotherapeutic group: Adrenergic, inhalants. Selective beta-2 adrenoreceptor agonists

ATC code: R03AC02

Salbutamol is a selective beta₂-adrenoceptor agonist. At therapeutic doses it acts on the beta₂-adrenoceptors of bronchial muscle providing short acting (4-6 hour) bronchodilation with a fast onset (within 5 minutes) in reversible airways obstruction.

Special Patient Populations

Children < 4 years of age

Pediatric clinical studies conducted at the recommended dose (SB020001, SB030001, SB030002), in patients < 4 years with bronchospasm associated with reversible obstructive airways disease, show that SALBIN INHALER has a safety profile comparable to that in children ≥ 4 years, adolescents and adults.

5.2 Pharmacokinetic properties

Salbutamol administered intravenously has a half-life of 4 to 6 hours and is cleared partly renally and partly by metabolism to the inactive 4'-O-sulfate (phenolic sulfate) which is also excreted primarily in the urine. The faeces are a minor route of excretion.

After administration by the inhaled route between 10 and 20% of the dose reaches the lower airways. The remainder is retained in the delivery system or is deposited in the oropharynx from where it is swallowed. The fraction deposited in the airways is absorbed into the pulmonary tissues and circulation, but is not metabolized by the lung. On reaching the systemic circulation it becomes accessible to hepatic metabolism and is excreted, primarily in the urine, as unchanged drug and as the phenolic sulfate.

The swallowed portion of an inhaled dose is absorbed from the gastrointestinal tract and undergoes considerable first-pass metabolism to the phenolic sulfate. Both unchanged drug and conjugate are excreted primarily in the urine. Most of a dose of salbutamol given intravenously, orally or by inhalation is excreted within 72 hours. Salbutamol is bound to plasma proteins to the extent of 10%.

5.3 Preclinical safety data

In common with other potent selective beta₂-agonists, salbutamol has been shown to be teratogenic in mice when given subcutaneously. In a reproductive study, 9.3% of foetuses were found to have cleft palate at 2.5mg/kg dose. In rats, treatment at the levels of 0.5, 2.32, 10.75 and 50 mg/kg/day orally throughout pregnancy resulted in no significant foetal abnormalities. The only toxic effect was an increase in neonatal mortality at the highest dose level as the result of lack of maternal care. Reproductive studies in the rabbit at doses of 50 mg/kg/day orally (i.e. much higher than the normal human dose) have shown foetuses with treatment related changes; these included open eyelids (ablepharia), secondary palate clefts (palatoschisis), changes in ossification of the frontal bones of the cranium (cranioschisis) and limb flexure.

Reformulation of the SALBIN INHALER has not altered the known toxicological profile of salbutamol.

In an oral fertility and general reproductive performance study in rats at doses of 2 and 50 mg/kg/day, with the exception of a reduction in number of weanlings surviving to day 21 post partum at 50 mg/kg/day, there were no adverse effects on fertility, embryofoetal development, litter size, birth weight or growth rate.

The non-CFC propellant, HFA 134a, has been shown to have no toxic effect at very high vapor concentrations, far in excess of those likely to be experienced by patients, in a wide range of animal species exposed daily for periods of two years.

6. Pharmaceutical particulars

6.1 List of excipients

1, 1,1,2-Tetrafluoroethane(Norflurane)

Oleic Acid

Ethanol

6.2 Incompatibilities

None reported.

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store below 30° C.

Protect from frost and direct sunlight.

As with most inhaled medications in aerosol canisters, the therapeutic effect of this medication may decrease when the canister is cold.

This canister contains a pressurized liquid. Do not expose to temperatures higher than 50°C. The canister should not be broken, punctured or burnt, even when empty.

Replace the mouthpiece cover firmly and snap it into position.

6.5 Nature and contents of container

An inhaler comprising an aluminum alloy can sealed with a metering valve, actuator and dust cap. Each canister contains 200 metered actuations providing 100 micrograms of salbutamol (as Salbutamol Sulfate BP).

6.6 Special precautions for disposal and other handling

The aerosol spray is inhaled through the mouth into the lungs. After shaking the inhaler, the mouthpiece is placed in the mouth and the lips closed around it. The actuator is depressed to release a spray, which must coincide with inspiration of breath.

For detailed instructions for use refer to the Patient Information Leaflet in every pack.

7. Marketing authorization holder

AMB HK ENTERPRISES (PVT) LTD

7.1

Product License Holder & Manufacturer: M/s Heilongjiang Province Fulekang Pharmaceutical Co., Ltd., No. 3 Yingbin St. NiuJia Industrial Park South Industrial City Harbin China.

8. Marketing authorization number(s)

115774

9. Date of first authorization/renewal of the authorization

Date of first authorization: 19TH JULY, 2024

Date of latest renewal: Not applicable

10. Date of revision of the text

Not applicable

Company Contact Details

AMB HK ENTERPRISES (PVT) LTD

Address

2ND FLOOR PLAZA 60, COMMERCIAL BLOCK-K, PHASE-1, DHA, LAHORE

Telephone

042 37207408

E-mail

Amb-hk-enterprises@hotmail.com